

Table 2. Combined estimates from random-effects meta-analyses for residential air pollution and respiratory infections during early life (up to 36 months).^a

	Crude ^b			Adjusted ^c		
	OR (95% CI)	I ²	p-Value	OR (95% CI)	I ²	p-Value
Pneumonia						
NO ₂	1.25 (1.04, 1.50)*	37.1	0.112	1.30 (1.02, 1.65)*	52.9	0.024
NO _x	1.23 (1.06, 1.41)*	22.2	0.239	1.26 (1.04, 1.52)*	44.0	0.066
PM _{2.5}	2.13 (0.82, 5.49)	79.7	0.000	2.58 (0.91, 7.27)	81.7	0.000
PM _{2.5} absorbance	1.78 (1.30, 2.43)*	0	0.734	1.99 (1.44, 2.75)*	0	0.663
PM ₁₀	1.55 (1.03, 2.34)*	29.2	0.205	1.76 (1.00, 3.09)*	51.2	0.051
Coarse PM	1.23 (1.02, 1.47)*	0	0.626	1.24 (1.03, 1.50)*	0	0.579
Traffic, nearest street	1.08 (1.03, 1.14)*	0	0.997	1.09 (1.03, 1.15)*	0	0.969
Traffic, major streets	1.19 (1.08, 1.31)*	0	0.979	1.21 (1.09, 1.34)*	0	0.843
Otitis media						
NO ₂	1.08* (1.01, 1.15)	4.8	0.395	1.09 (1.02, 1.16)*	0	0.515
NO _x	1.04 (0.98, 1.10)	0.5	0.430	1.05 (0.98, 1.12)	0	0.458
PM _{2.5}	1.02 (0.71, 1.45)	55.5	0.047	1.06 (0.75, 1.49)	47.9	0.088
PM _{2.5} absorbance	1.05 (0.80, 1.37)	46.7	0.095	1.08 (0.83, 1.39)	39.9	0.139
PM ₁₀	0.98 (0.83, 1.17)	11.6	0.341	0.98 (0.84, 1.14)	0	0.539
Coarse PM	0.96 (0.87, 1.06)	0	0.608	0.97 (0.88, 1.08)	0	0.805
Traffic, nearest street	0.98 (0.94, 1.03)	1.4	0.385	0.98 (0.93, 1.02)	0	0.497
Traffic, major streets	1.00 (0.91, 1.09)	0	0.462	0.99 (0.89, 1.10)	18.2	0.300
Croup						
NO ₂	0.92 (0.80, 1.07)	0	0.884	0.96 (0.83, 1.12)	0	0.909
NO _x	0.96 (0.83, 1.10)	0	0.895	0.99 (0.86, 1.14)	0	0.936
PM _{2.5}	0.83 (0.58, 1.19)	0	0.760	0.90 (0.63, 1.30)	0	0.703
PM _{2.5} absorbance	0.95 (0.66, 1.37)	5.0	0.368	1.03 (0.72, 1.47)	0	0.554
PM ₁₀	0.89 (0.70, 1.13)	0	0.586	0.92 (0.72, 1.18)	0	0.595
Coarse PM	0.95 (0.80, 1.12)	0	0.551	0.97 (0.82, 1.15)	0	0.787
Traffic, nearest street	0.98 (0.93, 1.04)	0	0.926	0.99 (0.93, 1.05)	0	0.853
Traffic, major streets	0.97 (0.86, 1.09)	0	0.734	0.98 (0.87, 1.11)	0	0.901

Associations are presented for the following increments in exposure: 10 µg/m³ for NO₂, 20 µg/m³ for NO_x, 5 µg/m³ for PM_{2.5}, 1 unit for PM_{2.5} absorbance, 10 µg/m³ for PM₁₀, 5 µg/m³ for coarse PM, 5,000 vehicles/day for traffic intensity on the nearest street; and 4,000 vehicle-km/day for traffic load on major roads within a 100-m buffer; associations with traffic intensity and traffic load were additionally adjusted for background NO₂ concentrations.

^aOutcomes assessed up to 12 months (INMA Valencia), 14 months (INMA Gipuzkoa, INMA Sabadell), 15 months (GASPII), 18 months (INMA Asturias), 24 months (BAMSE, GINIplus, LISAPLUS, PIAMA), and 36 months (MAAS). ^bCrude models were adjusted for sex and municipality (BAMSE). ^cAdjusted models included municipality (BAMSE), sex, older siblings, breastfeeding at 6 months, atopy of either parent, any child-care reported during follow-up, maternal smoking during pregnancy, any environmental tobacco smoke in the child's home reported during follow-up, visible mold or dampness in the home, use of gas stove, birth season, parental socioeconomic status (low, medium, high), and intervention (GINIplus, MAAS, PIAMA). **p* < 0.05.

Table 3. Adjusted combined estimates for air pollution exposure at the birth address and respiratory infection by year of life [OR (95% CI)].

	Pneumonia ^a (<i>n</i> = 12,891)	Otitis media ^b (<i>n</i> = 8,722)	Croup ^c (<i>n</i> = 9,101)
Respiratory infections during the first year ^d of life			
NO ₂	1.47* (1.15, 1.89)	1.19* (1.07, 1.33)	1.05 (0.83, 1.32)
NO _x	1.45* (1.21, 1.75)	1.09 (0.98, 1.22)	1.10 (0.90, 1.36)
PM _{2.5}	4.06* (1.93, 8.57)	1.21 (0.64, 2.28)	1.15 (0.67, 1.97)
PM _{2.5} absorbance	2.71* (1.68, 4.37)	1.32 (0.99, 1.75)	1.04 (0.59, 1.83)
PM ₁₀	1.77* (1.18, 2.67)	1.24 (0.76, 2.02)	1.07 (0.75, 1.53)
Coarse PM	1.46* (1.11, 1.92)	1.16 (0.80, 1.70)	1.02 (0.80, 1.30)
Traffic, nearest street	1.14* (1.07, 1.22)	0.99 (0.94, 1.04)	1.03 (0.94, 1.13)
Traffic, major streets	1.31* (1.15, 1.50)	1.03 (0.93, 1.14)	1.00 (0.81, 1.24)
Respiratory infections during the second year ^e of life			
NO ₂	1.40* (1.04, 1.88)	1.07 (0.96, 1.20)	0.92 (0.78, 1.09)
NO _x	1.29* (1.07, 1.55)	1.02 (0.89, 1.17)	0.92 (0.78, 1.08)
PM _{2.5}	2.65 (0.63, 11.2)	1.06 (0.64, 1.74)	0.76 (0.51, 1.15)
PM _{2.5} absorbance	1.90 (0.93, 3.87)	1.20 (0.80, 1.79)	0.89 (0.59, 1.35)
PM ₁₀	1.42 (0.99, 2.03)	1.00 (0.84, 1.19)	0.83 (0.63, 1.09)
Coarse PM	1.24 (0.98, 1.56)	1.00 (0.89, 1.13)	0.89 (0.73, 1.08)
Traffic, nearest street	1.05 (0.98, 1.13)	0.96 (0.90, 1.03)	0.93 (0.81, 1.07)
Traffic, major streets	1.10 (0.90, 1.34)	0.96 (0.83, 1.10)	1.00 (0.88, 1.14)

Associations are presented for the following increments in exposure: 10 µg/m³ for NO₂, 20 µg/m³ for NO_x, 5 µg/m³ for PM_{2.5}, 1 unit for PM_{2.5} absorbance, 10 µg/m³ for PM₁₀, 5 µg/m³ for coarse PM, 5,000 vehicles/day for traffic intensity on the nearest street; and 4,000 vehicle-km/day for traffic load on major roads within a 100-m buffer; associations with traffic intensity and traffic load were additionally adjusted for background NO₂ concentrations.

^aBased on four studies: BAMSE, GINI/LISA North, GINI/LISA South, PIAMA. ^bBased on 3 studies: BAMSE, LISAPLUS North, LISAPLUS South, PIAMA. ^cBased on three studies: BAMSE, GINI/LISA North, GINI/LISA South. ^dDefined as 0–12 months. ^eDefined as 13–24 months. Models were adjusted for municipality (BAMSE), sex, older siblings, breastfeeding at 6 months, atopy of either parent, any child-care reported during follow-up, maternal smoking during pregnancy, any environmental tobacco smoke in the child's home reported during follow-up, visible mold or dampness in the home, use of gas stove, birth season, parental socioeconomic status (low, medium, high), and intervention (GINIplus, PIAMA). **p* < 0.05.

diagnosis is complicated by the fact that not all infections present with acute symptoms severe enough to warrant a physician visit, in contrast with pneumonia, which routinely presents with a high fever and/or difficulty breathing (Edmond et al. 2012).

Conclusion

Our meta-analysis of 10 European birth cohorts found consistent evidence for an association between traffic-related air pollution and pneumonia, and some evidence to suggest an association with otitis media. Policies aimed at reducing air pollution may be successful in reducing the overall burden of pneumonia in early childhood.

CORRECTION

The value for “Traffic load on major streets within 100-m buffer” for GINI/LISA South in Table 1 was incorrect in the manuscript originally published online. It has been corrected here.

REFERENCES

- Aguilera I, Pedersen M, Garcia-Esteban R, Ballester F, Basterrechea M, Esplugues A, et al. 2013. Early-life exposure to outdoor air pollution and respiratory health, ear infections, and eczema in infants from the INMA study. *Environ Health Perspect* 121:387–392; doi:10.1289/ehp.1205281.
- Bateson TF, Schwartz J. 2008. Children's response to air pollutants. *J Toxicol Environ Health A* 71(3):238–243.
- Beelen R, Hoek G, Vienneau D, Eeftens M, Dimakopoulou M, Pedeli X, et al. 2013. Development of NO₂ and NO_x land use regression models for estimating air pollution exposure in 36 study areas in Europe—the ESCAPE project. *Atmos Environ* 72:10–23.
- Black RE, Cousens S, Johnson HL, Lawn JE, Rudan I, Bassani DG, et al. 2010. Global, regional, and national causes of child mortality in 2008: a systematic analysis. *Lancet* 375:1969–1987.
- Brauer M, Gehring U, Brunekreef B, de Jongste J, Gerritsen J, Rovers M, et al. 2006. Traffic-related air pollution and otitis media. *Environ Health Perspect* 114:1414–1418; doi:10.1289/ehp.9089.
- Brunekreef B, Smit J, de Jongste J, Neijens H, Gerritsen J, Postma D, et al. 2002. The Prevention and Incidence of Asthma and Mite Allergy (PIAMA) birth cohort study: design and first results. *Pediatr Allergy Immunol* 13:55–60.
- Cesaroni G, Porta D, Badaloni C, Stafoggia M, Eeftens M, Meliefste K, et al. 2012. Nitrogen dioxide levels estimated from land use regression models several years apart and association with mortality in a large cohort study. *Environ Health* 11:48; doi:10.1186/1476-069X-11-48.
- Clarke RW, Antonini JM, Hemenway DR, Frank R, Kleeberger SR, Jakab GJ. 2000. Inhaled particle-bound sulfate: effects on pulmonary inflammatory responses and alveolar macrophage function. *Inhal Toxicol* 12(3):169–186.
- Custovic A, Simpson BM, Murray CS, Lowe L, Woodcock A; NAC Manchester Asthma and Allergy Study Group. 2002. The National Asthma Campaign Manchester Asthma and Allergy Study. *Pediatr Allergy Immunol* 13(suppl 15):32–37.
- Cyrys J, Eeftens M, Heinrich J, Ampe C, Armengaud A, Beelen R, et al. 2012. Variation of NO₂ and NO_x concentrations between and within 36 European study areas: results from the ESCAPE study. *Atmos Environ* 62: 374–390.
- da Costa JL, Navarro A, Neves JB, Martin M. 2004. Household wood and charcoal smoke increases risk of otitis media in childhood in Maputo. *Int J Epidemiol* 33(3):573–578.
- DerSimonian R, Laird N. 1986. Meta-analysis in clinical trials. *Control Clin Trials* 7(3):177–188.
- D'Souza-Vazirani D, Minkovitz CS, Strobino DM. 2005. Validity